

# Bimodal HD-KFM and Resiscope Atomic Force Microscopy characterization of bidimensional materials and solar cells.

Paco Martinez<sup>1</sup> and Louis Pacheco<sup>2</sup>

<sup>1</sup>ScienTec Iberica, Rufino Sanchez 83, Las Rozas, España

<sup>2</sup>Concept Scientific Instruments, 17 Rue des Andes, Les Ulis, France


[f.martinez@scientec.es](mailto:f.martinez@scientec.es)

## Abstract

Over the past 30 years, Atomic Force Microscopy has evolved from a microscope to measure just the surface topography to a wide variety of measurement modes that provides a way to characterize other atomic interactions or physical properties like magnetic field, electric field, nanoscale dissipation processes, thermal conductivity, electrical conductivity, resistance, surface potential, piezoresponse, Young modulus,... Electrical nanocharacterization with AFM has emerged as a powerful tool to map electrical properties at the nanoscale, like surface potential (work function) and conductivity. However, traditional setups in AFM make difficult to obtain accurate and repeateable results over several types of samples.

In this article we will show the capabilities of two new developed AFM modes: High Definition Kelvin Force Microscopy (HD-KFM) and (Soft)Resiscope that overcome the intrinsic difficulties of electrical nanocharacterization with AFM. This two techniques have been applied on a wide variety of substrates: bidimensional materials, like graphene or molibdene disulfide, organic solar cells and nanoparticles providing high stability, sensitivity and lateral resolution.

## Figures



a) HD-KFM image on Graphene b) HD-KFM image on Molibdene disulfide

- 1.G. Binnig, C.F. Quate, Ch. Gerber, Phys. Rev. Lett. 56, 930 (1986).
- 2.Houzé F, Meyer R, Schneegans O, Boyer L.. Appl Phys Lett. 1996;69:1975.
- 3.D.W. Abraham, et al, J. Vac. Sci. Technol. B 9,703 (1991)
- 4.T.R. Albrecht, P. Grütter, D. Horne, D. Rugar, J. Appl. Phys. 69, 668 (1991).
- 5.T.R.Rodriguez and R.Garcia App. Phys. Lett. 84(3):449-451
6. J. Colchero, A. Gil, A.M. Baró, Phys. Rev. B 64, 245403 (2001)