

# Acoustic THz graphene plasmons revealed by photocurrent nanoscopy

Pablo Alonso-González<sup>1,2</sup>, Alexey Y. Nikitin<sup>1,3</sup>, Yuanda Gao<sup>4</sup>, Achim Woessner<sup>5</sup>, Mark B. Lundeberg<sup>5</sup>, Alessandro Principi<sup>6</sup>, Nicolò Forcellini<sup>7</sup>, Wenjing Yan<sup>1</sup>, Saül Vélez<sup>1</sup>, Andreas. J. Huber<sup>8</sup>, Kenji Watanabe<sup>9</sup>, Takashi Taniguchi<sup>9</sup>, Luis E. Hueso<sup>1,3</sup>, Marco Polini<sup>10</sup>, James Hone<sup>4</sup>, Frank H. L. Koppens<sup>5,11</sup>, and Rainer Hillenbrand<sup>3,12</sup>

<sup>1</sup>CIC nanoGUNE, E-20018, Donostia-San Sebastián, Spain,

<sup>2</sup>Departamento de Física, Universidad de Oviedo, Oviedo 33007, Spain,

<sup>3</sup>IKERBASQUE, Basque Foundation for Science, 48011 Bilbao, Spain,

<sup>4</sup>Department of Mechanical Engineering, Columbia University, New York, NY 10027, USA,

<sup>5</sup>ICFO-Institut de Ciències Fotoniques, The Barcelona Institute of Science and Technology, 08860 Castelldefels (Barcelona), Spain,

<sup>6</sup>Radboud University, Institute for Molecules and Materials, NL-6525 AJ Nijmegen, The Netherlands,

<sup>7</sup>Department of Physics, Imperial College London, London SW7 2AZ, United Kingdom,

<sup>8</sup>Neaspec GmbH, Martinsried, Germany,

<sup>9</sup>National Institute for Materials Science, 1-1 Namiki, Tsukuba 305-0044, Japan,

<sup>10</sup>Istituto Italiano di Tecnologia, Graphene labs, Via Morego 30 I-16163 Genova, Italy,

<sup>11</sup>ICREA – Institució Catalana de Recerca i Estudis Avançats, Barcelona, Spain,

<sup>12</sup>CIC NanoGUNE and EHU/UPV, E-20018, Donostia-San Sebastian, Spain

pabloalonso@uniovi.es

## Abstract:

The interaction of terahertz (THz) radiation with graphene has a vast application potential in many technologies, including imaging, communications, sensing, or photo-detection, among others. Recently, it has been shown that the excitation of localized THz plasmons in graphene can strongly enhance light-matter interactions, opening the door to more efficient optoelectronic devices. Here, we will present on the first visualization of propagating graphene plasmons (GPs) at THz frequencies, which can also be controlled by metallic (split) gates. Intriguingly, due to the coupling of GPs with the metal gate underneath we observe a linearization of the plasmon dispersion (thus revealing acoustic plasmons), which comes along with an extreme confinement of the plasmon fields [1]. These extraordinary GPs properties are very promising for sensing and communication technologies. To map the THz GPs, we introduce nanoscale-resolved THz photocurrent nanoscopy as a novel tool for studying fundamental and applied aspects of local THz photocurrent generation with a resolution of 25 nm, nearly 4 orders of magnitude below the diffraction limit.

## References

- [1] P. Alonso-González, A. Y. Nikitin, Y. Gao, A. Woessner, M. B. Lundeberg, A. Principi, N. Forcellini, W. Yan, S. Vélez, A. J. Huber, K. Watanabe, T. Taniguchi, L. E. Hueso, M. Polini, J. Hone, F. H. L. Koppens, and R. Hillenbrand, *Nature Nanotechnology*, doi:10.1038/nnano.2016.185 (2016).